

Heterocyclic terpenes: linear furano- and pyrroloterpenoids

Yonghong Liu,^{*a} Si Zhang^a and Pedro J. M. Abreu^b

Received (in Cambridge, UK) 30th March 2006

First published as an Advance Article on the web 6th June 2006

DOI: 10.1039/b604586c

Covering: January 1970 to December 2005

The emphasis of this review is on linear furano- and pyrroloterpenoids, together with their relevant biological activities, source organisms and country of origin. First total syntheses that lead to the revision of structures or stereochemistries have been included, and 206 references are cited.

1	Introduction
2	Reviews
3	Furanoterpenoids

3.1	Furanosesquiterpenoids
3.2	Furanoditerpenoids
3.3	Furanosesterterpenoids
3.4	Furanotriterpenoids
4	Pyrroloterpenoids
4.1	Pyrrolosesquiterpenoids
4.2	Pyrrolosesterterpenoids and norsesterterpenoids
5	Acknowledgements
6	References

^aGuangdong Key Laboratory of Marine Materia Medica, South China Sea Institute of Oceanology, Chinese Academy of Sciences, Guangzhou, 510-301, China. E-mail: yhliu6@yahoo.com.cn; Fax: +86-20-84451672; Tel: +86-20-89023105

^bCQFB/REQUIMTE, Departamento de Química, Faculdade de Ciências e Tecnologia, Universidade Nova de Lisboa, Quinta da Torre, 2829-516, Caparica, Portugal

Yonghong Liu obtained his BS and MS degrees from the Changchun College of Traditional Chinese Medicine, China. He then went to Korea, and received his PhD degree from Pusan National University, where he studied the isolation, structure elucidation and structure–activity relationships of biologically active marine natural products under the guidance of Professor Jee H. Jung. He undertook post-doctoral research with Nobutoshi Murakami at Osaka University, Japan, and Pedro Abreu at the New University of Lisbon, Portugal, before returning to China to take a position as a senior scientist at the South China Sea Institute of Oceanology, Chinese Academy of Sciences. His research interests are the isolation and structural elucidation of biologically active secondary metabolites from marine organisms.

Si Zhang received his BS degree from Qingdao Ocean University, followed by an MS degree from the South China Sea Institute of Oceanology, Chinese Academy of Sciences. After receiving his PhD degree from Guangdong University of Traditional Chinese Medicine, China, he is now a professor at the South China Sea Institute of Oceanology, a director of the National (Sanya) Key Research Field-station, and a director of Guangdong Key Laboratory of Marine Materia Medica. His research interests involve the properties, structure and activity of components from marine organisms. He has accomplished the development of two kinds of new marine drugs, and the preclinical study of a new marine drug expected to become a national key drug.

Pedro Abreu graduated in chemistry at the Free University of Brussels (Belgium) in 1975, and returned to Portugal that year to work in the chemical industry. In 1978 he joined the Chemistry Department of the Faculty of Science and Technology (FCT), New University of Lisbon, where he received his PhD in Organic Chemistry in 1987. After a post-doctoral appointment with Professor Takao Yokota at the Faculty of Agriculture, University of Tokyo (1987), he returned to FCT to pursue his teaching and research activities as Professor of Natural Products Chemistry. In the last decade the research interests have been focused on the isolation, characterization, and biological evaluation of natural products from African and Portuguese medicinal plants, and algae of the Portuguese coast.



Yonghong Liu



Si Zhang



Pedro Abreu

1 Introduction

This review covers the literature from 1970 to 2005 and describes 326 compounds from 206 articles. We show the structures of novel compounds, and previously reported compounds where there has been a structural revision or a newly established stereochemistry. Previously reported compounds for which first syntheses or new bioactivities are described have been referenced, but separate structures have not generally been shown.

2 Reviews



A number of reviews have dealt with natural sesquiterpenoids,¹ diterpenoids,² sesterterpenoids³ and triterpenoids.⁴ General reviews include those of marine natural products.^{5,6} Where appropriate, references to other reviews have been included in the following sections.

3 Furanoterpenoids


3.1 Furanosesquiterpenoids

3.1.1 Monofuranosesquiterpenoids.

3.1.1.1 Dendrolasin-type furanosesquiterpenoids. Dendrolasin **1** is a very well-known compound, and four syntheses have been reported.^{7–10} An investigation of the sponge *Dysidea herbacea*, collected on the Great Barrier Reef, has led to the identification of sesquiterpenoid **2**.¹¹ A furanosesquiterpene **3** has been isolated from the marine sponge *Dysidea fragilis* (Mont.) from Brittany.¹² An isomer of dehydrodendrolasin **4** has been isolated from the Australian nudibranch *Ceratosoma brevicaudatum*, collected around the jetty at Rapid Bay, South Australia.¹³ The absolute configuration of (+)-myoporone **5**, a furanosesquiterpenoid from *Myoporum montanum*,¹⁴ was determined by enantioselective synthesis.¹⁵ Dictyodendrillins A–C **6–8** were isolated from a South Australian *Dictyodendrillia* species.¹⁶ Dictyodendrillin B **7** was prepared by a concise route in 43% overall yield.¹⁷




The *R* configuration of the hydroxyl group in 4-hydroxymyoporone (athanagrandione) **9**¹⁸ has been established by an interrelationship with (+)-eremoacetal.¹⁹ The first total synthesis of the stress factor (+)-hydroxymyoporone **9** has been established and its absolute configuration reassigned.²⁰ The furanoterpene isodehydrodendrolasin **10** has been obtained from a tropical marine sponge *Euryspongia deliculata*, collected at Heron Island on the Great Barrier Reef, Australia.²¹ Two homofarnesane lactones, **11** and **12**, and farnesane lactone **13** have been found in the aerial parts of *Gochnatia glutinosa*. This is the first time that homofarnesanes have been isolated from plants, although several compounds of this type have been obtained from insects.²²




Four sesquiterpenoids **14–17** that are related biosynthetically to ipomeamarone, the well-known sweet potato phytoalexin, have been isolated from the root tissue of sweet potato (*Ipomoea batatas*) infected with the fungus *Cerutocystis fimbriata*.²³ Japanese workers have obtained another phytoalexin, **18**,¹⁴ from this system

in the same way. An isodehydrodendrolasin derivative **19** has been obtained from nudibranchs of the Cantabrian Sea belonging to the *Hypselodoris* genus, which use this compound as a deterrent against predators.²⁴




Six furanosesquiterpenes of the dendrolasin type, **20–25**, have been isolated from a cytotoxic extract of the deep ocean tunicate *Ritterella rete*,²⁵ whilst known members of this sesquiterpene group have been found in the Patagonian nudibranch *Tyrinna nobilis*.²⁶ The sesquiterpene **26** has been isolated from *Eremophila forrestii*, a medicinal plant in the arid regions of Western Australia.²⁷




3.1.1.2 Other types of furanosesquiterpenoids. Continuing their studies of Australian soft corals, Bowden *et al.* have isolated twelve structurally related furanosesquiterpenes, **27–38**, from *Simularia capillosa*.²⁸ Compounds **27**, **28**, and **31** were extremely acid-


sensitive and decomposed on storage. The product **28** was also isolated from *Simularia firma*.²⁸ The dehydrovernopolanthofuran **39** has been obtained from *Vernonia polyanthes*.²⁹ Practical syntheses of the anti-inflammatory furanosesquiterpene **36**, a metabolite of *Simularia* sp.,²⁸ employ a strategy based on the Claisen rearrangement.³⁰



The furanosesquiterpenoid **40** was isolated from the Mediterranean alcyonacean *Alcyonium pulmatum*.³¹ A furanosesquiterpene **41** has been obtained from the soft coral *Lobophytum catalai*, which was collected from the Andaman and Nicobar Islands.³² Actinolide A **42** is a sesquiterpenic lactone, found in an extract of the stems of *Actinodaphne lancifolia*.³³










Anthecotulide **43** has been isolated from *Anthemis cotula*.³⁴ The isolation and structure determination of two furanosesquiterpenes, **44** and **45**, from the soft coral *Simularia lochmodes* have been described.³⁵ Another *Simularia* sp. contains the furanosesquiterpene **44**.³⁶ A sesquiterpene named algoafuran **46** has been found in an extract of the nudibranch *Leminda millecra*, collected in Algoa Bay, South Africa.³⁷ Siphonodictidine **47** has been synthesized for the first time in a concise and regiocontrolled manner in an overall yield of 25.7%.³⁸ Siphonodictidine **47** is the secondary metabolite of an Indo-Pacific sponge *Siphonodictyon* sp.,³⁹ which burrows




into living coral. Among the members of this class discovered so far,^{40,41} siphonodictidine is remarkable for its biological action,³⁹ as it inhibits growth of the coral and appears to be deployed by the sponge to kill coral polyps in its immediate vicinity. Apart from this specific function, which is essential for the survival of the sponge, siphonodictidine exhibits significant antifungal and antimicrobial activity against Gram-positive and Gram-negative bacteria.

The sesquiterpenes **48–50** have been isolated from *Arctotis arctotoides*.⁴² The sesquiterpenoid **51** has been isolated from the root tissue of sweet potato (*Ipomoea batatas*) infected with the fungus *Cerutocystis fimbriata*.²³ Compound **52**, which possesses a sesquiterpenoid skeleton of the caleprunane type, has been isolated from *Calea prunifolia*.⁴³ The structures of cyclonerodiol oxide and epicyclonerodiol oxide, which were isolated from the fungus *Trichoderma polysporum*, have been shown to be **53** and **54**, respectively.⁴⁴








The sesquiterpene **55**, related to davanone, has been isolated from *Artemisia reptans*.⁴⁵ The new nerolidol derivative arteincultone **56** has been isolated from *Artemisia inculta*.⁴⁶ The essential oil of *Artemisia pallens* (davana oil) contains more than 50 components, one of which was identified as the nerolidol derivative


57.⁴⁷ A peroxynerolidol derivative, fercoperol **58**, has been found in *Ferula communis* ssp. *communis*.⁴⁸ A cyclic sesquiterpene peroxysemiketal, **59**, has been isolated from *Artemisia maritima*.⁴⁹



The davanone-type peroxides epiarteincultone **60** and spiroarteincultone **61** have been extracted from *Artemisia abrotanum*. These two metabolites were also prepared synthetically by singlet and triplet oxygen oxidation respectively, and **61** showed schizonticidal activity against *Plasmodium falciparum*.⁵⁰ The essential oil of *Santolina oblongifolia* contains the *cis*- and two *trans*-isomers of **62**.⁵¹ The structures of (+)-myomontanone **63** and isomyomontanone **64**, which are furanosesquiterpenoids from *Myoporum montanum*, were shown to be the aldol condensation products of (+)-myoporone **5**, which is also present in this plant.¹⁴




The furanosesquiterpene merrekentrone D **65** has been found in the roots and rootstocks of *Merremia kentrokaulus*.⁵² Four furanosesquiterpene β -hydroxy ketones previously isolated from *Myoporum* sp.⁵³ have now been assigned complete stereostructures from NMR and X-ray data. The compounds are (5*R*,7*R*,9*S*)-9-hydroxy-5,9-dihydromyomontanone **66**, (5*S*,7*R*,9*R*)-9-hydroxy-5,9-dihydromyomontanone **67**, (5*R*,7*R*,9*R*)-9-hydroxy-5,9-dihydromyomontanone **68** and (4*S*,7*S*,8*R*)-4-hydroxy-10,11-didehydro-4,8-dihydromyodesmone **69**.







Plakortone G **70** is a highly cytotoxic sesquiterpene, isolated from a Caribbean sponge of the *Plakortis* genus.⁵⁴ Two furanosesquiterpenoid ketones, dihydrocrassifolone **71** and crassifolone **72**, were isolated from the wood essential oil of *Myoporum crassifolium* from New Caledonia.⁵⁵ The furanosesquiterpene laevidiene **73** has been isolated from *Cadlina iaevis* from the Cantabrian Sea (Atlantic Ocean).⁵⁶ The structure **74** has been assigned to a furanosesquiterpenoid obtained from the extract of the aerial parts of *Ixiolaena leptolepis*.⁵⁷ Two new furanosesquiterpene alcohols, pelseneeriol 1 **75** and pelseneeriol 2 **76**, have been isolated from the porostome nudibranch *Doriopsilla pelseneeri*, collected off the Portuguese coast (Atlantic Ocean).⁵⁸ An approach known to give regiospecific and stereospecific reactions has been applied to the first enantioselective synthesis of the natural (–)-microcionin 2 **77**, an example of a molecule containing a *trans* relationship between the methyl groups.⁵⁹ Microcionin was first

isolated from the sponge *Microcionia toxystila*,⁶⁰ and later from the dorid nudibranch molluscs (*Dendrodoris grandiflora*) growing in contact with them.⁶¹ A second group has also isolated this furanosesquiterpene from another species of dorid nudibranch molluscs, *Cadlina luteomarginata*.⁶²






Tetraenol **78** has been isolated from the relict Chinese plant *Tetraena mongolica*.⁶³ The incubation of small colonies of *Simularia capillosa*, freshly collected from Magnetic Island (Australia), with labeled mevalonolactone afforded the furanoquinol **79**, containing the label in the terpenoid portion of the sesquiterpene hydroquinone; this confirmed how mevalonate is incorporated into terpenoids *in vivo* in this soft coral.⁶⁴ The sesquiterpenoid derivative **80** has been obtained from the roots of *Ferula feruloides*.⁶⁵ The first total synthesis of cristic acid **81** has been reported.⁶⁶ This bioactive sesquiterpene had previously been isolated from the fruiting bodies of the higher mushroom *Albatrellus cristatus*.⁶⁷ An asymmetric synthesis of (*R*)-sesquilavandulol has been achieved.⁶⁸ *Ferula feruloides* also contains the farnesane derivatives **82** and **83**.^{69,70}



Four sesquiterpene coumarins **84**–**87**, showing nitric oxide production inhibitory effects, have been found in *Ferula fukanensis*.^{71,72} The two dihydrofurochromone sesquiterpenes **88** and **89** have been obtained from the roots of *Ferula ferulaeoides*.⁷³



Five sesquiterpene phenylpropanoid derivatives, fukanedones A–E **90–94**, were isolated from the roots of *Ferula fukanensis*.



These compounds inhibited NO production and inducible NO synthase gene expression by a murine macrophage-like cell line activated by lipopolysaccharide and recombinant mouse interferon- γ .⁷⁴


An efficient and stereodefined process is described for the first preparation of a prenyl-benzoylfuranone-type sesquiterpenoid, (\pm)-3-(2,4-dihydroxybenzoyl)-4,5-dimethyl-5-(4,8-dimethylnona-3(*E*),7(*E*)-dien-1-yl)tetrahydro-2-furanone **83**.⁷⁰ The synthetic strategy is based on the stereoselective construction of the key quaternary carbon center by nucleophilic addition of organometallic reagents to functionalized ketoamides derived from dihydroxyacetone dimer.⁷⁵ Pallidones B–J **95–103** have been isolated from the EtOAc extracts of the roots of *Ferula pallida* (Steud.) Korovin (Umbelliferae), which grows in Bulgan Somon of Hovd City, Mongolia.^{76,77} Merrekentrones A **104** and B **105** have been found in the roots and rootstocks of *Merremia kentrokaulus*.⁵²


3.1.2 Bisfuranosesquiterpenoids. A variety of farnesyl/nerolidyl sesquiterpenoids from various plant sources have been identified. These include the lasiospermone derivative **106** from *Ursinia nana*,⁷⁸ and brachymerolide **107** and brachymeral **108** from *Brachymeris montana*,⁷⁹ one of the woody South African genera of the Anthemideae. The structure of a bisbutenolide, a likely precursor of the acetylenic sesquiterpene freelingyne, obtained from a species of the genus *Eremophila*, has been determined as **109**.⁸⁰ A nerolidol derivative, **110**, has been obtained from *Ursinia sericeae*.⁸¹ The sesquiterpene bisbutenolides **111** and **112** have been isolated from *Eremophila homoplastica*.²⁷

The work described in a detailed paper by Russell and Sutherland has confirmed the absolute configuration of (–)-ngaione **113**, and furthermore it has been conclusively demonstrated that ipomeamarone, a phytoalexin from the sweet potato, is in fact (+)-ngaione **114**.⁸² On this basis, it is suggested that the use of the name ipomeamarone should be discontinued. Biosynthetic studies have shown that the three oxygen atoms in (+)-ngaione **114** (= ipomeamarone) are derived from molecular oxygen, not from water nor from the hydroxyl group of farnesol itself.⁸³ The absolute configuration of (+)-ipomeamarone **114** has also been independently determined.⁸⁴ A furanosesquiterpene, merrekentrone C **115**, has been found in the roots and rootstocks of *Merremia kentrokaulus*.⁵² A sesquiterpenoids that are related biosynthetically to ipomeamarone which had been infected with the fungus *Cerutocystis fimbriata*, has the structure **116**.²³ A study of the roots of *Rehmannia glutinosa* led to the isolation of a sesquiterpene **117**.⁸⁵ Another species of this genus, *Ferula ferulioides*, contains farnesane derivatives **118** and **119**.¹¹

3.1.3 Trisfuranosesquiterpenoids. A sesquiterpenoid **120** related biosynthetically to ipomeamarone **114**, the well-known sweet potato phytoalexin, has been isolated from root tissue of sweet potato (*Ipomoea batatas*) infected with the fungus *Cerutocystis fimbriata*.²³ Japanese workers have obtained⁸⁶ another phytoalexin **121**, from this system in the same way.


The sesquiterpene chilene B **122**, a trimer of 2-methyl-3(*2H*)-furanone, has been obtained from the red alga *Laurencia chilensis*, and its structure determined by X-ray analysis.⁸⁷ The synthesis of the difuranosesquiterpene athanasin **123** has enabled its absolute configuration to be established.⁸⁸ Stereochemical studies






on the biosynthesis of homoterpenes derived from nerolidol and geranyllinalool from higher plants, together with mechanistic, phylogenetic, and ecological aspects, have been reported.⁸⁹

3.1.4 Norfuranosesquiterpenoids. The Hawaiian sponge *Spongia oceania* contains pokepola ester **124**, a diester of phosphoric acid in which the alcohols are 5-methylhexan-1-ol and homoserine. The amine of homoserine forms an amide with a trinorfuranosesquiterpenic acid.⁹⁰ The norsesquiterpene **125** has been found in an extract of the aerial parts of *Artemisia inculta*, collected in three different locations in Egypt.⁹¹ The sesquiterpenes **126** and **127** have been obtained from *Athanasia pinnata*.⁹² The aerial part of *Calea crocinervosa* contains the trisnorsesquiterpene lactone crocinervolide **128**.⁹³ Compound **129**, which possesses the caleprunane skeleton, has been isolated from *Calea prunifolia*.⁴³ The norsesquiterpene **130** has been isolated from *Artemisia herba-alba*.⁹⁴




3.2 Furanoditerpenoids

3.2.1 Monofuranoditerpenoids. The furan furosolidagonone **131** is among the constituents of *Solidago drummondii*.⁹⁵ Microglossic acid **132** (from *Microglossa zeylanica* collected in Sri Lanka)⁹⁶ and 17-(acetoxymethyl)thymifodioic acid **132** (from *Baccharis thymifolia*)⁹⁷ are further examples. It has been suggested that the occurrence of hydroxylated geranylgeraniol derivative **134** with




a butenolide end-group may have taxonomic significance in the classification of the tribe Astereae (Compositae).⁹⁸


The eleanolone diterpenoid **135** was amongst a series obtained from the alga *Bifurcaria bifurcata*.⁹⁹ Two diterpenes, **136** and **137**, isolated from a Western Australian species of *Spongia*, were obtained as 3 : 1 mixtures of enantiomers.¹⁰⁰ The brown alga *Bifurcaria bifurcata*, collected from Brittany on the Atlantic coast, contained two diterpenoids, bifurcane **138** (which inhibited the development of fertilized sea urchin eggs), and epoxyeleanolactone **139**.¹⁰¹



The furanoditerpene **140** was obtained from the Australian sponge *Thorectandra choanoides*.¹⁰² The geranylgeraniol derivative **141** has been isolated from a Brazilian medicinal plant, erva-santa, *Baccharis ocracea* (Asteraceae).¹⁰³ Incanic acid **142** from *Conyza incana* possesses an *E* alkene geometry.¹⁰⁴ Inconyzic acid **143** was obtained from *Conyza incana*.¹⁰⁵








Compound **144** was obtained from specimens of the brown alga *Bifurcaria bifurcata* from two locations in Brittany.¹⁰⁶ Cacospongionolide C **145** was obtained from the Adriatic sponge *Fasciospongia caverrzosa*.¹⁰⁷ A number of acyclic diterpenoids possess a terminal furan ring, exemplified by conyzaleucolide A **146** obtained from *Conyza hypoleuca* (Compositae).¹⁰⁸




Marine organisms continue to yield oxygenated diterpenoids, of which thuridillin **147**, from the mollusc *Thuridilla hopei*, is an example.¹⁰⁹ Linear diterpenoids have been considered as chemotaxonomic markers in the genus *Cystoseira* (brown algae).¹¹⁰ Halitunal **148** has been isolated from the marine alga *Halimeda tuna*,¹¹¹ and two acyclic furanoditerpene compounds, saurufuran A **149** and B **150**, have been obtained from the root of *Saururus chinensis*. Saurufuran A is a peroxisome proliferator-activated receptor γ (PPAR γ) agonist, however, saurufuran B weakly activated PPAR γ .¹¹² The geranyl-geraniol derivative **151** has been isolated from *Diplostephium meyenii* (Asteraceae).¹¹³


Lingulatusin **152** and tuxpanolide **153** are geranyl-geraniol derivatives reported from *Aster ligulatus*¹¹⁴ and *Perymenium hintonii*,¹¹⁵ respectively. An acyclic diterpene- γ -lactone, hanliuine I **154**, has been obtained from *Salix matsudan*.¹¹⁶ The lactone naviculide **155** has been obtained from the liverwort *Porella navicularis*,¹¹⁷ whilst some geranyl-linalool glycosides, the capsiansides,




have been reported from *Capsicum* plants.¹¹⁸ The lactone **156** has been isolated from *Melampodium leucanthum*.¹¹⁹



Hanliuin III **157** is a lactone obtained from *Salix matsudana*.¹²⁰ Cacospongine A **158** was isolated from a Philippine sponge, *Cacospongia* sp.¹²¹ Examination of the resinous exudate of a *Madia* species has yielded madiaol **159**.¹²² Examination of the sponge *Jaspis splendens* afforded (+)-subersin **160**, which is an inhibitor of 15-lipoxygenase.¹²³




3.2.2 Bisfuranoditerpenoids. Caucanolides A–F **161**–**166** have been isolated from the gorgonian octocoral *Pseudopterogorgia bipinnata* collected in the south-western Caribbean Sea near Colombia. These compounds showed *in vitro* antiplasmodial activity against the malaria parasite *Plasmodium falciparum*.¹²⁴ A Korean collection of a sponge of the genus *Sarcotragus* was found to contain two bisfuranoditerpenoids, sarcotins K **167** and L **168**, which displayed weak cytotoxicity.¹²⁵




3.3 Furanosesterterpenoids

3.3.1 C₂₁ furanoterpenoids. Many C₂₁ furanosesterterpenes contain a furan ring at both ends of the molecule, such as nitenin **169** and dihydronitenin **170** from *Spongia nitens*,¹²⁶ and anhydrofurospongine-1 **171**¹²⁷ and furospongine-1 **172**¹²⁸ from *Spongia officinalis* and *Hippospongia communis*, respectively. The compounds **173** and **174** have been detected in a species of the genera *Spongia* and *Leiosella*, respectively,¹²⁹ whilst the γ -lactone furospongolide **175**¹³⁰ has been found in *Dysidea herbacea*.




The sponge *Cacospongia scalaris* has provided a number of linear C₂₁ difurans, exemplified by cacospongione A **176**, cacospongione B **177**, and cacospongione C **178**.¹³¹ The unsaturated lactone **179**, kurospongine, having ichthyotoxic and feeding-deterrent properties, has been isolated from a marine sponge *Spongia* sp.¹³² Its absolute stereochemistry was established by applying Horeau's method to an alcohol derived from a Grignard reaction on the unsaturated lactone. The untenospongins A **180** and B **181**, obtained from a *Hippospongia* species,¹³³ possess

coronary vasodilating properties. The sponge *Carteriospongia flabellifera*, collected from the Great Barrier Reef, contains 12,13-dehydrofurospingin 1 **182**.¹³⁴




Degraded linear sesterterpenes have been isolated from sponges of the genera *Spongia*, *Carteriospongia* and *Hippospongia*. For example, untenospongins C **183** has been obtained from an Okinawan *Hippospongia* sp.¹³⁵ Isonitenin **184** is an additional C₂₁ difuranoterpene from a Spanish collection of *Spongia officinalis*.¹³⁶ The degradation of tetronic acids with alkaline hydrogen peroxide, to mimic the formation of these compounds, has been reported.¹³⁷


Ent-kurospingin **185** was isolated from a Korean *Sarcotragus* species.¹³⁸ A *Hippospongia* sp. from Goa, India, contained *ent*-untenospongins A **186**.¹³⁹



A C₂₁ difuranoterpene, **187**, from a Mediterranean specimen of *Spongia virgultosa*, was found to be the enantiomer of (–)-untenospongins B from a Japanese *Hippospongia* sp.^{133,140} Comparison of spectral data suggested that the structure of tetrahydrofurospingin-1 from an Australian *Spongia* sp.¹²⁹ should be revised from **188** to **187**.¹⁴⁰ The absolute configuration of the C₂₁ difuranoterpene **189**, isolated from *Fasciospongia cavernosa* from the Arabian Sea,¹⁴¹ has been determined by synthesis of its enantiomer.¹⁴²




A specimen of *Spongia officinalis* from La Caleta, Cádiz, Spain, contained two additional minor C₂₁ furanoterpenes, the weakly cytotoxic furospingin-5 **190** and cyclofurospingin 2 **191**.¹⁴³ A C₂₁ norsesterterpenoid originally reported to have two conjugated double bonds, **192**,¹²⁹ has been revised to **193** on the basis of more complete spectroscopic data obtained from a sample isolated from an Australian specimen of *Spirastrella papilosa*.¹⁴⁴ The absolute stereochemistry was determined by degradation, and the name (–)-isotetrahydrofurospingin-1 proposed for this bisfuranoterpene.




The suggestion that ircinin 3 **194** and ircinin 4 **195** may arise by cleavage of the open forms of the tetronic acids¹⁴⁵ has been confirmed by the occurrence of the corresponding C₂₁ carboxylic acids **196–200** in *Ircinia dendroides*.¹⁴⁶ A synthesis of ircinin 4 **195**, a metabolite of *Ircinia oros*,¹⁴⁵ employed a palladium-catalyzed reaction as the key step.¹⁴⁷


3.3.2 C₂₂ norsesterterpenoids. A C₂₂ γ -lactone, furodendin **201**, has been reported in the sponge *Phyllospongia dendyi*.¹⁴⁸ A furanoterpene, designated dehydrofurodendin **202**, was isolated from two different species of Madagascan sponges of the genus *Lendenfeldia*, both of which seem to be as yet undescribed.¹⁴⁹ Dehydrofurodendin **202** has been found to be a potent inhibitor of HIV-1 RT-associated DNA polymerase activity. It should be noted that **202** differs from **201** only in the additional double bond at C-16–C-17, which probably results from oxidation occurring at a late stage in its biosynthesis.






Hippospongins E **203** and F **204**, from *Hippospongia* sp. of southern Australian waters, are truncated furanosesterterpenes incorporating diaminoethane.¹⁵⁰ A Korean collection of a sponge of the genus *Sarcotragus* was found to contain sarcotins I **205** and J **206**. These compounds have a unique C₂₂ trinorsesterterpene skeleton, and might be degradation products of the corresponding sesterterpenes.¹²⁵









A moderately cytotoxic norsesterterpenoid, sarcotin N **207**, has been isolated from a Korean *Sarcotragus* species, and, in the same study, the previously reported sarcotin I **205** was found to have the 21*R* configuration.¹³⁸ Two trinorsesterterpenoid lactams, the sarcotragins A **208** and B **209**, have been isolated from a *Sarcotragus* species from Jaeju Island (Korea).¹⁵¹












3.3.3 C₂₄ norsesterterpenoids. Konakhin **210**, from a Senegalese sponge, represents a degradation product of the corresponding tetronic acid, and is possibly derived from fasciculatin.¹⁵² The open form of the tetronic acid is a β -keto acid, and on decarboxylation it generates the enol as the substrate for chloroperoxidase. It is worth noticing that the rate-determining step in many electrophilic halogenations is the formation of the enol, and it is therefore possible that the decarboxylation and halogenation steps are biosynthetically linked. Two similar bisfurans, **211** and **212**, have been obtained from a North Adriatic collection of *Ircinia oros*,¹⁵³ and were found to be toxic to *Artemia salina* shrimp. Three norsesterterpenoids, **213–215**, isolated from an Okinawa *Ircinia* species, were found to be moderately cytotoxic against the KB cell line.¹⁵⁴







A new C₂₄ furanoterpene, hippospongion D **216**, was isolated from a southern Australian collection of the marine sponge *Hippospongia* sp.¹⁵⁰ A moderately cytotoxic norsesterterpenoid, sarcotin O **217**, along with two pyrrolonorsesterterpenoids, sarcotrine F **218** and isosarcotrine F **219**, were isolated from a Korean *Sarcotragus* species.¹³⁸






3.3.4 C₂₅ non-conjugated tetronic acid furanosesterterpenoids. Palinurin **220** is a linear furanosesterterpene previously reported from the Mediterranean sponge *Ircinia variabilis*.¹⁵⁵ Ircinic acid **221** was isolated from marine sponge *Ircinia* sp. collected from Fiji and also found from a Truck lagoon sponge.¹⁵⁶ Palominin **222** is another geometrical isomer obtained¹⁵⁷ from a Caribbean *Ircinia* species. Like many of these compounds from sponges, it possesses ichthyotoxic properties and thus may have a role in protecting the sponge against predators.



Compound **223** has been reported as a constituent of *Psarnnocinia rugosa*.¹⁵⁸ The bisfurans spongionellin **224** and dehydrospongionellin have been described in extracts of a *Spongionella* species.¹⁵⁹ The okinonellins A **225** and B **226** are furanosesterterpenes from another *Spongionella* species, which inhibit cell-division in fertilized starfish eggs and again may play a role in marine chemical ecology.¹⁶⁰ Okinellin A **225**, isolated from the Okinawan marine sponge *Hippospongia* sp., is probably identical to hippospongion, and possesses antispasmodic activity.¹⁶¹ Both the 8*S*,21*S*,22*S*,23*R*- and 8*R*,21*S*,22*S*,23*R*-isomers of okinonellin B **226**, which is a cytotoxic and antispasmodic agent from *Spongionella* sp., have been synthesized, but neither has the same optical rotation as the natural product.¹⁶²








A tetronic acid with antimicrobial activity, **227**, has been isolated from an Australian sponge of the *Psammocinia* genus.¹⁶³ Isopalinurin **228**, which is a mild protein phosphatase inhibitor, was obtained from a South Australian *Dysidea* species.¹⁶⁴





Cometin A **229** contains two furan rings and is another product obtained from Australian sponges.¹⁶⁵ Three cytotoxic furanosesterterpenoids, sacotins A–C **230–232**, were reported from a specimen of *Sarcotragus* sp. collected at Cheju Island, Korea.¹⁶⁶ The cytotoxic sesterterpenoids *epi*-sacotin A **233**, and sacotins F **234** and M **235**, were isolated from same Korean *Sarcotragus* species.¹²⁵ *Epi*-sacotin F **236**¹³⁸ was found in two Korean *Sarcotragus* species.

Cytotoxic furanosesterterpenes palinurin **220**¹⁵⁵ and isopalinurin **228**¹⁶⁴ were re-isolated from a marine sponge *Psammocinia* sp. collected from Ulleung Island, Korea. The configuration at C-21 has not been previously investigated, and it has now been determined to be *R* on the basis of CD analysis.¹⁶⁷










3.3.5 C₂₅ conjugated tetronic acid furanosesterterpenoids. Marine organisms, particularly sponges, have continued to provide a source of linear sesterterpenoids. The terminal units often comprise either a furan or a tetronic acid moiety. Two isomeric linear difuranosesterterpenes, ircinin 1 **237** and ircinin 2 **238**, were obtained from the sponge *Ircinia oros*, and contain a second furan ring.¹⁶⁸ Quite a number of these compounds contain a tetronic acid moiety and show strong antibiotic activity against *Staphylococcus aureus*. These are exemplified by variabilin **239** from *Ircinia variabilis*,¹⁶⁹ its double-bond isomer strobilin **240** from *Ircinia strobilina*,¹⁷⁰ and fasciculatin **241** from *Ircinia fasciculata*.¹⁷¹






Compounds **242** and **243** occur in *Ircinia dendroides*.¹⁴⁶ Variabilin **239** and its double-bond isomers are quite widespread. As well as being obtained from sponges of the genus *Ircinia*, it has now been found in the New Zealand sponge *Sarcotragus*,¹⁷² whilst 22-deoxyvariabilin **244** has been found in a *Thorecta* species collected

off the Bay of Islands, New Zealand.¹⁷³ In an antimicrobial assay, 22-deoxyvariabilin inhibited the growth of *Staphylococcus aureus*, *Bacillus subtilis*, and *Candida albicans*. NMR studies of ircinins 1 **237** and 2 **238**, two difuranosesterterpenes isolated from a Fijian *Ircinia* sp., have been reported.¹⁵⁶ Two additional oxygenated furanosesterterpenes, **245** and **246**, have been described from an *Ircinia* species collected from Kaikoura, New Zealand.¹⁷⁴






Variabilin is readily available from *Ircinia*, *Psammocinia*, and *Sarcotragus* species, and because it shows promising activity as an antiviral agent, its chemistry has been explored.¹⁷⁵ The tetronic acid moiety acts as a sensitizer for the production of singlet oxygen, which then reacts with the furan to give a range of products including hydroxybutenolides. The 22-*O*-methyl derivative is stable, and although it retains antiviral activity, it was not possible to dissociate the cytotoxicity from the antiviral activity. The decomposition of these furans has stimulated further

studies¹⁷⁶ on the regioselective oxidation of 3-alkylfurans with singlet oxygen. (18*S*)-Variabilin **247** is a widespread member of this series, and its absolute stereochemistry and that of some related tetronic acids have been established.^{177,178} (18*R*)-Variabilin **248** has been isolated from the sponge *Ircinia felix* from Colombia.¹⁷⁹ Two sesterterpenoids, **249** and **250**, isolated from an Okinawa *Ircinia* species, were found to be moderately cytotoxic.¹⁵⁴





A re-investigation of the structure of strobilin, originally isolated from *Ircinia strobilina*,¹⁷⁰ revealed that it consisted of two geometrical isomers, **251** and **252**.¹⁸⁰ The structures of (18*S*)-ircinin-1 **237**,¹⁶⁸ (18*S*)-ircinin 2 **238**,¹⁶⁸ and strobilins **251** and **252** have been tentatively assigned.¹⁷⁸ Two additional geometrical isomers, (8*Z*,13*Z*,20*Z*)-strobilin **253** and (7*Z*,13*Z*,20*Z*)-felixin **255**, were partially characterized after isolation as their 22-*O*-acetates from Columbian specimens of *I. felix*, *I. strobilina* and *I. campana*.¹⁸¹ (8*Z*,13*Z*,18*S*,20*Z*)-Strobilin **254** and (7*E*,12*Z*,18*S*,20*Z*)-variabilin **256** from a Maltese specimen of *I. oros* were characterized as their corresponding 22-*O*-methyl derivatives.¹⁸² The first asymmetric synthesis of the marine furanosesterterpene tetronic acid 18*S*-variabilin **256**, from the sponge *Ircinia variabilis*, has been reported.¹⁸³ An efficient and stereodefined process is employed, featuring lipase-catalyzed asymmetric desymmetrization of two types of propanediol precursors incorporating the terpene skeleton.










Three new furanosesterterpenes, **257–259**, and the known cytotoxic compounds (7*E*,12*E*,18*R*,20*Z*)-variabilin, (8*E*,13*Z*,18*R*,20*Z*)-strobilin, (7*E*,13*Z*,18*R*,20*Z*)-felixinin [also known as (7*E*,13*Z*,18*R*,20*Z*)-variabilin], (8*Z*,13*Z*,18*R*,20*Z*)-strobilin, and (7*Z*,13*Z*,18*R*,20*Z*)-felixinin, were isolated from a Korean marine sponge, *Psammocinia* sp. These compounds were evaluated for cytotoxicity against a panel of five human tumor cell lines, and most of the compounds showed toxicity to SK-MEL-2.¹⁶⁷ A mixture of (8*E*,13*Z*,20*Z*)-strobilin and (7*E*,13*Z*,20*Z*)-felixinin displayed significant inhibition of DNA replication and a moderate antioxidant profile.¹⁸⁴





Eight cytotoxic furanosesterterpenoids, namely ircinins 1 and 2, sacotins D **260** and E **261**,¹⁶⁶ sacotins G **262** and H **263**,¹²⁵ (7*E*,12*E*,18*R*,20*Z*)-variabilin, and (8*E*,13*Z*,18*R*,20*Z*)-strobilin,¹³⁸ were reported from a specimen of *Sarcotragus* sp. collected off Cheju Island, Korea. 22-Deoxy-23-(hydroxymethyl)variabilin **264** was isolated from both the South African nudibranch *Hypselodoris capensis* and the sponge *Fasciospongia* sp. on which it was feeding.¹⁸⁵


3.3.6 C₂₅ monofuranosesterterpenoids. Simple furans are exemplified by furospongins 3 **265** and furospongins 4 **266**, identified from *Spongia officinalis*.¹⁴⁵ The sponge *Spongia idia* (*Leiosella idia*) from Pt Loma, San Diego, contained idiadione **267**.¹⁸⁶ Idiadione was found to be toxic to the sea star *Pisaster giganteus*, a brine shrimp *Artemia* sp., and the ectoproct *Membranipora memoranacea*, and




also immobilized the larvae of the red abalone *Haliotis rufescens* in sea-water. Idiadione **267** was synthesized from citronellal, establishing the stereogenic centre as *S*.¹⁸⁷




The sesterterpene **268**, obtained from the Caribbean sponge *Thorecta horridus*,¹⁸⁸ possessed a marked inflammatory activity, inducing the release of histamine and causing oedema in the paw of test animals. In contrast, thorectolide mono-acetate **269**, obtained from a New Caledonian marine sponge of the genus *Hyr-tios*,¹⁸⁹ possessed anti-inflammatory properties. The epoxyfuran carboxylic acid **270** has been isolated from a Western Australian *Spongia* species.¹⁰⁰ A cytotoxic sesterterpene, cacospongionolide D **271**, was isolated from *Fasciospongia cavernosa* from the Bay of Naples.¹⁹⁰ A specimen of *Spongia officinalis* from La Caleta, Cádiz, Spain, contained demethylfurospongins 4 **272**, which was weakly cytotoxic against the P-388 cell line.¹⁴³ A linear C₂₅ furanoterpene, furosposulin 1 **273**, has been isolated from *Ircinia spinosula*.¹⁹¹ The bicyclic lactone astakolactin **274** was obtained from specimens of *Cacospongia scalaris* collected from the Gulf of Astakos, Greece.¹⁹²




268




269




270




271



272




273




274

3.3.7 C₂₅ bisfuranosesterterpenoids. Luffarin R **275** has been isolated¹⁹³ from *Luffariella geometrica*. A cytotoxic sesterterpenoid, (6*Z*)-luffarin-V **276**, has been isolated from *Fasciospongia cavernosa* from the Bay of Naples.¹⁹⁰




275




276


3.3.8 C₂₅ trisfuranosesterterpenoids. Three C₂₅ furanoterpenes, hippospongins A–C **277–279**, were isolated from a southern Australian collection of the marine sponge *Hippospongia* sp. Hippospongin A **279** was found to be a mild antibiotic, inhibiting the growth of *Staphylococcus aureus*.¹⁵⁰



277




278




279


3.3.9 Salts of sesterterpenoids. Four unstable sulfate esters of furanosesterterpenes, **280–283**, were obtained from *Ircinia variabilis* and *I. oros* of the northern Adriatic Sea.¹⁹⁴ A Californian sponge of the family Halichondriidae contained sulfated sesterterpene halisulfate **3** **284**,¹⁹⁵ which possesses antimicrobial and anti-inflammatory properties. The 22-*O*-sulfates **285** and **286** of palinurin and fasciculatin, which were toxic to brine shrimp *Artemia salina*, were isolated from *Ircinia variabilis* and *I. fasciculata*, respectively.¹⁹⁶ A sodium salt of the pyrroloterpenoid sarcotragin A **208** was isolated from a *Sarcotragus* sp. collected from Jaeju Island in Korean waters.¹⁵¹ A sodium salt of trinorsesterterpene acid sarcotin I **205**,¹²⁵ and the sodium salts of four pyrroloterpenoids, sarcotrines E **287** and F **288** and isosarcotrines E **218** and F **219**,¹³⁸ were isolated from the Korean sponge *Sarcotragus* sp.




280



281




282



283


3.3.10 Esters of sesterterpenoids. (18*R*)-Variabilin-11-methyloctadecanoate ester **289** has been isolated from the sponge *Ircinia fefix* from Colombia.¹⁹⁷ The same sponge has yielded nine novel fatty acid esters, (7*E*,12*E*,18*R*,20*Z*)-variabilin




(5*Z*,9*Z*)-22-methyltricosadienoate, (7*E*,12*E*,18*R*,20*Z*)-variabilin (5*Z*,9*Z*)-tetracosadienoate, (7*E*,12*E*,18*R*,20*Z*)-variabilin hexadecanoate, (7*E*,12*E*,18*R*,20*Z*)-variabilin 10-methylhexadecanoate, (7*E*,12*E*,18*R*,20*Z*)-variabilin 15-methylhexadecanoate, (7*E*,12*E*,18*R*,20*Z*)-variabilin 14-methylhexadecanoate, (7*E*,12*E*,18*R*,20*Z*)-variabilin 9-octadecenoate, (7*E*,12*E*,18*R*,20*Z*)-variabilin octadecanoate, and (7*E*,12*E*,18*R*,20*Z*)-variabilin 2,11-dimethyloctadecanoate (**290–298**).¹⁹⁸


3.4 Furanotriterpenoids

3.4.1 C₃₀ terpenoids. A C₃₀ furanoterpene, furospinosulin **299**, has been isolated from *Ircinia spinosula*.¹⁹¹ Three triterpenoids, viz. hydroxymokupalide **300**, acetoxymokupalide **301**, and mokupalide **302**, have been obtained from a species of the genus *Megalopastas*.^{199,200} The structure of mokupalide **302** was confirmed by total synthesis.²⁰¹




(*Z,E,E*)-Cupaniopsin A, (*Z,E,Z*)-cupaniopsin A, the 29-methyl ester of (*Z,E,E*)-cupaniopsin A, (*Z,E,Z*)-cupaniopsin C, 22-hydroxy-(*Z,E,Z*)-cupaniopsin C, 23-hydroxy-(*Z,E,Z*)-cupaniopsin C, (*Z,E,Z*)-cupaniopsin A methyl ester, and (*Z,E,Z*)-cupaniopsin E (**303–310**) were isolated from the bark of *Cupaniopsis trigonocarpa* Radlk. & Guillaumin, *C. azantha* Radlk. and *C. phallacrocarpa* F. Adema (Sapindaceae). Five of them exhibited potent binding activity on the peroxisome proliferator-activated receptor, the most potent being (*Z,E,E*)-cupaniopsin A **303**. This is the first report of linear furanotriterpenes isolated from the plant kingdom.²⁰²





- 303** 16E $R_1 = \text{CH}_3$, $R_2 = \text{CH}_3$, $R_3 = \text{COOH}$, $R_4 = \text{H}$
304 16Z $R_1 = \text{CH}_3$, $R_2 = \text{CH}_3$, $R_3 = \text{COOH}$, $R_4 = \text{H}$
305 16E $R_1 = \text{CH}_3$, $R_2 = \text{CH}_3$, $R_3 = \text{COOH}$, $R_4 = \text{CH}_3$
306 16Z $R_1 = \text{CH}_3$, $R_2 = \text{CH}_3$, $R_3 = \text{CH}_3$, $R_4 = \text{H}$
307 16Z $R_1 = \text{CH}_2\text{OH}$, $R_2 = \text{CH}_3$, $R_3 = \text{CH}_3$, $R_4 = \text{H}$
308 16Z $R_1 = \text{CH}_3$, $R_2 = \text{CH}_2\text{OH}$, $R_3 = \text{CH}_3$, $R_4 = \text{H}$
309 16Z $R_1 = \text{CH}_3$, $R_2 = \text{CH}_3$, $R_3 = \text{COOH}$, $R_4 = \text{CH}_3$







3.4.2 C₃₁ terpenoids. It has been suggested that a C₃₁ difuran-oterpene, difurospinosulin **311** (isolated from *Ircinia spinosula*), could be derived from a C₃₅ linear furanoterpene by the loss of four carbon atoms.¹⁹¹ A C₃₁ furanoterpene alcohol sulfate, named ircinol sulfate **312**, has been isolated from a marine sponge, *Ircinia* sp., collected off New Caledonia.²⁰³ Ircinol sulfate is the second example of a C₃₁ furanoterpene encountered in marine sponges.

3.4.3 C₃₅ terpenoids. A linear C₃₅ furanoterpene, furospinosulin-3 **313**, has been isolated from *Ircinia spinosula*.¹⁹¹

4 Pyrroloterpenoids

4.1 Pyrrolsesquiterpenoids




Pyrrolostatin **314**, a lipid peroxidation inhibitor, was isolated from the culture of *Streptomyces chrestomyceticus* EC40.²⁰⁴ Pyrrolostatin inhibited lipid peroxidation in rat brain homogenate. A *Streptomyces* sp. (NPS008187) isolated from a marine sediment collected in Alaska was found to produce three pyrrolsesquiterpenes, glyciapyrroles A **315**, B **316**, and C **317**. Glyciapyrrole A




inhibited both colorectal adenocarcinoma HT-29 and melanoma B16-F10 tumor cell growth.²⁰⁵

4.2 Pyrrolsesquiterpenoids and norsesterterpenoids


The pyrrolsesquiterpenes sarcotrines A **318**, B **320**, C **322**, D **324**, E **287** and F **218**, *epi*-sarcotrines A **319**, B **321** and C **323**, and isosarcotrines E **288** and F **219**^{125,138} have been isolated from a marine sponge of the genus *Sarcotragus*. Two trinorsesterterpenoid lactams, sarcotragins A **208** and B **209**, with structures similar to those of the sarcotrines, have been isolated from another *Sarcotragus* sponge collected from Jaegu Island in Korean waters.¹⁵¹ It is quite feasible that sarcotrines and sarcotragins share the same biogenetic precursor, and that the latter are the degradation products of the former. Two microbial metabolites of palinurin **220**¹⁵⁵ (re-isolated from a Red Sea sponge, *Ircinia echinata*), palinurines A **325** and B **326**, are produced by the fungus *Cunninghamella* sp. NRRL 5695.²⁰⁶ The possible transformation of furans to amides (*viz.* dehydro-3-enepyrrolidin-2-one and dehydro-3-enepyrrolidin-5-one) through a biomimetic reaction is certain to have applications in synthetic chemistry.






318 (21S), R =


319 (21R), R =




320 (21S), R =




321 (21R), R =




322 (21S), R =



323 (21R), R =



324 R =



325 R₁ = H₂, R₂ = O

326 R₁ = O, R₂ = H₂

5 Acknowledgements

We are grateful for a grant from the Hundred Talents Project of the Chinese Academy of Sciences.

6 References

- B. M. Fraga, *Nat. Prod. Rep.*, 2005, **22**, 465, and earlier reviews cited therein.
- J. R. Hanson, *Nat. Prod. Rep.*, 2005, **22**, 594, and earlier reviews cited therein.
- J. R. Hanson, *Nat. Prod. Rep.*, 1996, **13**, 529, and earlier reviews cited therein.
- J. D. Connolly and R. A. Hill, *Nat. Prod. Rep.*, 2005, **22**, 487, and earlier reviews cited therein.
- D. J. Faulkner, *Nat. Prod. Rep.*, 2002, **19**, 1, and earlier reviews cited therein.
- J. W. Blunt, B. R. Copp, M. H. G. Munro, P. T. Northcote and M. R. Prinsep, *Nat. Prod. Rep.*, 2005, **22**, 15, and earlier reviews cited therein.
- S. P. Tanis, *Tetrahedron Lett.*, 1982, **23**, 3115.
- S. Araki and Y. Butsugan, *Chem. Lett.*, 1982, 177.
- E. Lee, Y. H. Park and S. K. Park, *Tetrahedron Lett.*, 1982, **23**, 2671.
- D. K. Barma, A. Kundu, R. Baati, C. Mioskowski and J. R. Falck, *Org. Lett.*, 2002, **4**, 1387.
- R. W. Dunlop, R. Kazlauskas, G. March, P. T. Murphy and R. J. Wells, *Aust. J. Chem.*, 1982, **35**, 95.
- G. Guella, A. Guerriero, P. Traldi and F. Pietra, *Tetrahedron Lett.*, 1983, **24**, 3897.
- M. B. Ksebati and F. J. Schmitz, *J. Nat. Prod.*, 1988, **51**, 857.
- P. L. Metra and M. D. Sutherland, *Tetrahedron Lett.*, 1983, **24**, 1749.
- R. C. Anand and V. Singh, *Tetrahedron*, 1993, **49**, 6515.
- N. H. Tran, J. N. A. Hooper and R. J. Capon, *Aust. J. Chem.*, 1995, **48**, 1757.
- K. Gerlach and H. M. R. Hoffmann, *Synlett*, 1998, 682.
- L. T. Burka, L. Kuhnert, B. J. Wilson and T. M. Harris, *Tetrahedron Lett.*, 1974, **15**, 4017.
- E. Dimiitriadis and R. A. Massy-Westropp, *Phytochemistry*, 1984, **23**, 1325.
- L. F. Tietze, C. Wegner and C. Wulff, *Chem.-Eur. J.*, 1999, **5**, 2885.
- R. J. Clark, M. J. Garson, I. M. Brereton and J. A. Kennedy, *J. Nat. Prod.*, 1999, **62**, 915.
- M. I. Ybarra, C. A. N. Catalan, J. Diaz and W. Herz, *Phytochemistry*, 1994, **35**, 1585.
- J. A. Schneider, J. Lee, Y. Naya, K. Nakanishi, K. Oba and I. Uritani, *Phytochemistry*, 1984, **23**, 759.
- A. Fontana, C. Avila, E. Martinez, J. Ortea, E. Trivellone and G. Cimino, *J. Chem. Ecol.*, 1993, **19**, 339.
- L. A. Lenis, M. J. Ferreiro, C. Debitus, C. Jiménez, E. Quiñoa and R. Riguera, *Tetrahedron*, 1998, **54**, 5385.
- A. Fontana, C. Muniain and G. Cimino, *J. Nat. Prod.*, 1998, **61**, 1027.
- Y. M. Syah, E. M. Ghisalberti, B. W. Skelton and A. H. White, *J. Nat. Prod.*, 1997, **60**, 49.
- B. F. Bowden, J. C. Coll, E. D. de Silva, M. S. L. de Costa, P. J. Djura, M. Mahendran and D. M. Tapiolas, *Aust. J. Chem.*, 1983, **36**, 371.
- F. Bohlmann, C. Zdero, R. M. King and H. Robinson, *Phytochemistry*, 1983, **22**, 2863.
- D. H. Williams and D. J. Faulkner, *Tetrahedron*, 1996, **52**, 4245.
- G. Cimino, S. de Rosa, S. de Stefano and G. Sodano, *J. Nat. Prod.*, 1984, **47**, 877.
- A. S. R. Anjaneyulu, N. S. K. Rao and M. J. R. V. Venugopal, *Indian J. Chem., Sect. B: Org. Chem. Incl. Med. Chem.*, 1996, **35**, 1001.
- M. R. Kim, H. J. Jung, B. S. Min, S. R. Oh, C. S. Kim, K. S. Ahn, W. S. Kang and H. K. Lee, *Phytochemistry*, 2002, **59**, 861.
- R. N. Baruah, F. Bohlmann and R. M. King, *Planta Med.*, 1985, 531.
- S. K. Park and P. J. Scheuer, *J. Korean Chem. Soc.*, 1994, **38**, 749.
- S. Mizobuchi, K. Kon-ya, K. Adachi, M. Sakai and W. Miki, *Fish Sci.*, 1994, **60**, 345.
- K. L. McPhail, M. T. Davies-Coleman and J. Starmer, *J. Nat. Prod.*, 2001, **64**, 1183.
- C. W. Jefford, J.-C. Rossier, J. Boukouvalas, A. W. Sledeski and P.-Z. Huang, *J. Nat. Prod.*, 2004, **67**, 1383.
- B. Sullivan, D. J. Faulkner and L. Webb, *Science*, 1983, **221**, 1175.
- J. E. Hochlowski and D. J. Faulkner, *Tetrahedron Lett.*, 1981, **22**, 271.
- D. W. Knight and D. C. Rustidge, *J. Chem. Soc., Perkin Trans. 1*, 1981, 679.
- F. Tschritzis, J. Jakupovic and F. Bohlmann, *Phytochemistry*, 1990, **29**, 195.
- V. Castro, J. Jakupovic and F. Bohlmann, *J. Nat. Prod.*, 1984, **47**, 802.
- T. Fujita, Y. Takaishi, Y. Takeda, T. Fujiyama and T. Nishi, *Chem. Pharm. Bull.*, 1984, **32**, 4419.
- J. A. Marco, J. F. Sanz-Cervera, F. Sancenon, M. Arno and J. Valles-Xirau, *Phytochemistry*, 1994, **37**, 1095.
- S. M. Khafagy, M. A. Al-Yahya, J. Ziesche and F. Bohlmann, *Phytochemistry*, 1983, **22**, 1821.
- L. N. Misra, A. Chandra and R. S. Thakur, *Phytochemistry*, 1991, **30**, 549.
- M. Miski, T. J. Mabry and F. Bohlmann, *J. Nat. Prod.*, 1986, **49**, 916.
- G. Ruecker, E. Breitmaier, R. Mayer and D. Manns, *Arch. Pharm. (Weinheim, Ger.)*, 1987, **320**, 437.
- G. Ruecker, D. Manns and S. Wilbert, *Arch. Pharm. (Weinheim, Ger.)*, 1993, **326**, 457.
- J. De Pascual-Teresa, S. Vicente, M. S. Gonzalez and I. S. Bellido, *Phytochemistry*, 1983, **22**, 2235.
- K. Jenett-Siems, K. Siems, L. Witte and E. Eich, *J. Nat. Prod.*, 2001, **64**, 1471.

- 53 R. M. Carman, P. N. Handley, R. Kadirvelraj, W. T. Robinson and M. D. Sutherland, *Aust. J. Chem.*, 1999, **52**, 727.
- 54 D. J. Gochfeld and M. T. Hamann, *J. Nat. Prod.*, 2001, **64**, 1477.
- 55 C. Menut, P. Cabalion, E. Hnawia, H. Agnani, J. Waikiedre and A. Fruchier, *Flavour Fragrance J.*, 2005, **20**, 621.
- 56 A. Fontana, M. Gavagnin, E. Mollo, E. Trivellone, J. Orteat and G. Cimino, *Comp. Biochem. Physiol., B: Biochem. Mol. Biol.*, 1995, **111**, 283.
- 57 L. Lehmann, J. Jakupovic, F. Bohlmann, R. M. King and L. Haegi, *Phytochemistry*, 1988, **27**, 2994.
- 58 H. Gaspar, M. Gavagnin, G. Calado, F. Castelluccio, E. Mollo and G. Cimino, *Tetrahedron*, 2005, **61**, 11032.
- 59 S. Potvin and P. Canonne, *Tetrahedron: Asymmetry*, 1996, **7**, 2821.
- 60 G. Cimino, S. De Stefano, A. Guerriero and L. Minale, *Tetrahedron Lett.*, 1975, **43**, 3723.
- 61 G. Cimino, S. De Rosa, S. De Stefano, R. Morrone and G. Sodano, *Tetrahedron*, 1985, **41**, 1093.
- 62 J. Hellou, R. J. Andersen and J. E. Thompson, *Tetrahedron*, 1982, **38**, 1875.
- 63 Z. Y. Shao, Y. H. Zhang, J. N. Li, K. J. Jiang and B. K. Kuai, *Z. Naturforsch., C: Biosci.*, 2004, **59**, 181.
- 64 J. C. Coll, B. F. Bowden, D. M. Tapiolas, R. H. Willis, P. Djura, M. Streamer and L. Trott, *Tetrahedron*, 1985, **41**, 1085.
- 65 K. Kojima, K. Isaka, O. Purev, G. Jargalsaikhan, D. Suran, H. Mizukami and Y. Ogihara, *Chem. Pharm. Bull.*, 1998, **46**, 1781.
- 66 A. Fürstner and T. Gastner, *Org. Lett.*, 2000, **2**, 2467.
- 67 L. Zechlin, M. Wolf, W. Steglich and T. Anke, *Liebigs Ann. Chem.*, 1981, 2099.
- 68 S. Faure, J. D. Connolly, C. O. Fakunle and O. Piva, *Tetrahedron*, 2000, **56**, 9647.
- 69 K. Kojima, K. Isaka, O. Purev, G. Jargalsaikhan, D. Suran, H. Mizukami and Y. Ogihara, *Chem. Pharm. Bull.*, 1999, **47**, 690.
- 70 K. Kojima, K. Isaka, P. Ondognii, O. Zevgeegiiin, K. Davgiin, H. Mizukami and Y. Ogihara, *Chem. Pharm. Bull.*, 1999, **47**, 1145.
- 71 T. Motai, A. Daikonya and S. Kitanaka, *J. Nat. Prod.*, 2004, **67**, 432.
- 72 T. Motai and S. Kitanaka, *Chem. Pharm. Bull.*, 2004, **52**, 1215.
- 73 A. Nagatsu, K. Isaka, K. Kojima, P. Ondognii, O. Zevgeegiiin, P. Gombosurengiin, K. Davgiin, B. Irfan, C. M. Iqbal and Y. Ogihara, *Chem. Pharm. Bull.*, 2002, **50**, 675.
- 74 T. Motai and S. Kitanaka, *J. Nat. Prod.*, 2005, **68**, 365.
- 75 H. Yoda, K. Maruyama and K. Takabe, *Tetrahedron Lett.*, 2003, **44**, 1775.
- 76 B.-N. Su, Y. Takaishi, G. Honda, M. Itoh, Y. Takeda, O. K. Kodzhimatov and O. Ashurmetov, *J. Nat. Prod.*, 2000, **63**, 436.
- 77 B.-N. Su, Y. Takaishi, G. Honda, M. Itoh, Y. Takeda, O. K. Kodzhimatov and O. Ashurmetov, *J. Nat. Prod.*, 2000, **63**, 520.
- 78 F. Bohlmann and R. K. Gupta, *Phytochemistry*, 1982, **21**, 1309.
- 79 F. Bohlmann and C. Zdero, *Phytochemistry*, 1982, **21**, 1989.
- 80 K. A. Dastlik, P. G. Foster, E. L. Ghisalberti and P. R. Jefferies, *Phytochemistry*, 1989, **28**, 1425.
- 81 J. Jakupovic, U. Ganzer, P. Pritschow, L. Lehmann, F. Bohlmann and R. M. King, *Phytochemistry*, 1992, **31**, 863.
- 82 C. A. Russell and M. D. Sutherland, *Aust. J. Chem.*, 1982, **35**, 1881.
- 83 L. T. Burka and A. Thorsen, *Phytochemistry*, 1982, **21**, 869.
- 84 J. A. Schneider, K. Yoshihara and K. Nakanishi, *J. Chem. Soc., Chem. Commun.*, 1983, 352.
- 85 Y. Oshima, K. Tanaka and H. Hikino, *Phytochemistry*, 1993, **33**, 233.
- 86 I. Ito, N. Kato and I. Uritani, *Agric. Biol. Chem.*, 1984, **48**, 159.
- 87 A. San Martin, J. Rovirosa, C. Xu, H. S. M. Lu and J. Clardy, *Tetrahedron Lett.*, 1987, **28**, 6013.
- 88 G. Bojack and H. Bornowski, *Tetrahedron*, 1991, **47**, 9179.
- 89 A. Gaebler, W. Boland, U. Preiss and H. Simon, *Helv. Chim. Acta*, 1991, **74**, 1773.
- 90 R. S. Kalidindi, W. Y. Yoshida, J. A. Palermo, P. J. Sheuer and M. Kelly-Borges, *Tetrahedron Lett.*, 1994, **35**, 5579.
- 91 J. A. Marco, J. F. Saenz-Cervera, V. Garcia-Lliso, M. Guara and J. Valles-Xirau, *Phytochemistry*, 1997, **45**, 751.
- 92 C. Zdero, L. Lehmann and F. Bohlmann, *Phytochemistry*, 1991, **30**, 1161.
- 93 A. Ortega, J. C. Lopez and E. Maldonado, *Phytochemistry*, 1989, **28**, 2735.
- 94 J. A. Marco, J. F. Sanz-Cervera and G. Ocete, *J. Nat. Prod.*, 1994, **57**, 939.
- 95 F. Bohlmann, T. V. Chau-Thi, P. Singh and J. Jakupovic, *Planta Med.*, 1985, 487.
- 96 A. A. L. Gunatilaka, B. Dhanabalasingham, L. Paredes, J. Jakupovic, F. Bohlmann and N. K. B. Adikaram, *Phytochemistry*, 1987, **26**, 2408.
- 97 J. R. Saad, M. J. Pestchanker and O. S. Giordano, *Phytochemistry*, 1987, **26**, 3033.
- 98 C. Zdero, F. Bohlmann and H. M. Niemeyer, *Phytochemistry*, 1990, **29**, 1227.
- 99 L. Hougaard, U. Anthoni, C. Christopherson and P. H. Nielsen, *Phytochemistry*, 1991, **30**, 3049.
- 100 P. A. Searle and T. F. Molinski, *Tetrahedron*, 1994, **50**, 9893.
- 101 R. Valls, L. Piovetti, B. Banaigs, A. Archavlis and M. Pellegrini, *Phytochemistry*, 1995, **39**, 145.
- 102 S. Urban and R. J. Capon, *Aust. J. Chem.*, 1995, **48**, 1903.
- 103 E. P. Schenkel, A. M. Sobottka, P. P. Lankhorst and J. Schripsema, *Phytochemistry*, 1997, **44**, 755.
- 104 A. M. Galal, E. Abdel-Sattar, F. S. El-Feray, J. S. Massa, M. R. Meselhy, S. Kadota and T. Nambo, *Phytochemistry*, 1998, **48**, 159.
- 105 E. Abdel-Sattar, *Monatsh. Chem.*, 2001, **132**, 1095.
- 106 G. Culioli, M. Daoudi, V. Mesguiche, R. Valls and L. Piovetti, *Phytochemistry*, 1999, **52**, 1447.
- 107 S. De Rosa, R. Puliti, A. Crispino, A. De Giulio, C. De Sena, C. Jodice and C. A. Mattia, *Tetrahedron*, 1995, **51**, 10731.
- 108 C. Zdero, F. Bohlmann and G. M. Mungai, *Phytochemistry*, 1991, **30**, 575.
- 109 M. Gavagnin, A. Spinella, A. Crispino, R. De Almeida Epifanio, A. Marin and G. Cimino, *Gazz. Chim. Ital.*, 1993, **123**, 205.
- 110 R. Valls, L. Piovetti and A. Praud, *Hydrobiologia*, 1993, **260**, 549.
- 111 F. E. Koehn, S. P. Gunasekera, D. N. Niel and S. C. Cross, *Tetrahedron Lett.*, 1991, **32**, 169.
- 112 B. Y. Hwang, J. H. Lee, J. B. Nam, H. S. Kim, Y. S. Hong and J. J. Lee, *J. Nat. Prod.*, 2002, **65**, 616.
- 113 M. Bittner, A. Schuster and J. Jakupovic, *Phytochemistry*, 1991, **30**, 1329.
- 114 S. Yu, M. F. Wang, C.-T. Ho, C.-K. Chin, S.-W. Yang, G. A. Cordell, H. Lotter and H. Wagner, *Phytochemistry*, 1998, **49**, 609.
- 115 E. Maldonado, M. Bello, J. L. Villasenor and A. Ortega, *Phytochemistry*, 1998, **49**, 1115.
- 116 S. Zheng, J. Wang, J. Lu, T. Shen, L. Sun and X. Shen, *Planta Med.*, 2000, **66**, 487.
- 117 M. Toyota, F. Nagashima and Y. Asakawa, *Phytochemistry*, 1989, **28**, 1661.
- 118 S. Yahara, Y. Izumitani and T. Nohara, *Tennen Yuki Kagobutsu Toronkai Koen Yoshishu*, 1988, **30**, 173.
- 119 L. Quijano, F. R. Fronczek and N. H. Fischer, *Phytochemistry*, 1985, **24**, 1747.
- 120 J. X. Wang, S. Z. Zheng, L. P. Sun, X. W. Shen and Y. Li, *J. Chin. Chem. Soc. (Taipei, Taiwan)*, 2002, **49**, 437.
- 121 D. Tasdemir, M. K. Harper, G. P. Concepcion, G. C. Mangalindan, E. Hajdu and C. M. Ireland, *Tetrahedron*, 2000, **56**, 9025.
- 122 E. Wollenweber, M. Dorr, M. Dorsam, A. El-Hamed Hassan, A. A. Ahmed, M. F. Hegazy and K. P. Zeller, *Z. Naturforsch., C: Biosci.*, 2003, **58**, 153.
- 123 J. Carroll, E. N. Jonsson, R. Ebel, M. S. Hartman, T. R. Holman and P. Crews, *J. Org. Chem.*, 2001, **66**, 6847.
- 124 C. A. Ospina, A. D. Rodriguez, J. A. Sanchez, E. Ortega-Barria, T. L. Capson and M. S. Mayer, *J. Nat. Prod.*, 2005, **68**, 1519.
- 125 Y. Liu, J. Hong, C.-O. Lee, K. S. Im, N. D. Kim, J. S. Choi and J. H. Jung, *J. Nat. Prod.*, 2002, **65**, 1307.
- 126 E. Fattorusso, L. Minale, G. Sodano and E. Trivellone, *Tetrahedron*, 1971, **27**, 3909.
- 127 G. Cimino, S. de Stefano, L. Minale and E. Fattorusso, *Tetrahedron*, 1972, **28**, 267.
- 128 G. Cimino, S. de Stefano, L. Minale and E. Fattorusso, *Tetrahedron*, 1971, **27**, 4673.
- 129 R. J. Capon, E. L. Ghisalberti and P. R. Jefferies, *Experientia*, 1982, **38**, 1444.
- 130 Y. Kashman and M. Zviely, *Experientia*, 1980, **36**, 1179.
- 131 G. Guella, P. Amande and F. Pietra, *Helv. Chim. Acta*, 1986, **69**, 726.
- 132 J. Tanaka and T. Higa, *Tetrahedron*, 1988, **44**, 2805.
- 133 A. Umeyama, N. Shoji, S. Aihara, Y. Ohizumi and J. Kobayashi, *Aust. J. Chem.*, 1989, **42**, 459.
- 134 F. J. Schmitz and J. C. Chang, *J. Nat. Prod.*, 1988, **51**, 745.
- 135 J. Kobayashi, H. Shinonega, H. Shigemori and T. Sasaki, *Chem. Pharm. Bull.*, 1993, **41**, 381.
- 136 L. A. Lenis, L. Nunez, C. Jimenez and R. Riguera, *Nat. Prod. Lett.*, 1996, **8**, 15.

- 137 G. Guella, I. Mancini, I. N'Diaye and F. Pietra, *Tetrahedron Lett.*, 1991, **32**, 6415.
- 138 Y. Liu, T. A. Mansoor, J. Hong, C.-O. Lee, C. J. Sim, K. S. Im, N. D. Kim and J. H. Jung, *J. Nat. Prod.*, 2003, **66**, 1451.
- 139 Y. W. Guo and E. Trivellone, *Chin. Chem. Lett.*, 2000, **11**, 327.
- 140 A. Fontana, L. Albarella, G. Scognamiglio, M. Uriz and G. Cimino, *J. Nat. Prod.*, 1996, **59**, 869.
- 141 M. Kobayashi, R. Chavakula, O. Murata and N. S. Sarma, *Chem. Pharm. Bull.*, 1992, **40**, 599.
- 142 T. Bando and K. Shishido, *Chem. Commun.*, 1996, 1357.
- 143 L. Garrido, E. Zubia, M. J. Ortega and J. Salvá, *J. Nat. Prod.*, 1997, **60**, 794.
- 144 R. J. Capon, A. Jenkins, F. Rooney and E. L. Ghisalberty, *J. Nat. Prod.*, 2001, **64**, 638.
- 145 G. Cimino, S. De Stefano and L. Minale, *Tetrahedron*, 1972, **28**, 5983.
- 146 A. G. Gonzalez, M. L. Rodriguez and A. S. M. Barrientos, *J. Nat. Prod.*, 1983, **46**, 256.
- 147 A. Fürstner, T. Gastner and J. Rust, *Synlett*, 1999, 29.
- 148 R. Kazlauskas, P. T. Murphy and R. J. Wells, *Experientia*, 1980, **36**, 814.
- 149 L. Chill, A. Rudi, M. Aknin, S. Loya, A. Hizi and Y. Kashman, *Tetrahedron*, 2004, **60**, 10619.
- 150 S. J. Rochfort, D. Atkin, L. Hobbs and R. J. Capon, *J. Nat. Prod.*, 1996, **59**, 1024.
- 151 J. Shin, J. R. Rho, Y. Seo, H. S. Lee, K. W. Cho and C. J. Sim, *Tetrahedron Lett.*, 2001, **42**, 3005.
- 152 I. N'Diaye, G. Guella, I. Mancini, J. M. Kornprobst and F. Pietra, *J. Chem. Soc., Chem. Commun.*, 1991, 97.
- 153 A. De Giulio, S. De Rosa, G. D. Vincenzo, G. Strazzullo and N. Zavodnik, *J. Nat. Prod.*, 1990, **53**, 1503.
- 154 H. H. Issa, J. Tanaka and T. Higa, *J. Nat. Prod.*, 2003, **66**, 251.
- 155 G. Alfano, G. Cimino and S. de Stefano, *Experientia*, 1979, **35**, 1136.
- 156 L. V. Manes, P. Crews, M. B. Ksebati and F. J. Schmitz, *J. Nat. Prod.*, 1986, **49**, 787.
- 157 M. Ortiz-Garcia and A. D. Rodriguez, *Tetrahedron*, 1990, **46**, 1119.
- 158 V. Liokas, M. J. Garson and J. A. Carver, *Aust. J. Chem.*, 1989, **42**, 1805.
- 159 Y. Kato, N. Fusetani, S. Matsunaga and K. Hashimoto, *Chem. Lett.*, 1985, 1521.
- 160 Y. Kato, N. Fusetani, S. Matsunaga and K. Hashimoto, *Experientia*, 1986, **42**, 1299.
- 161 J. Kobayashi, Y. Ohizumi, H. Nakamura and Y. Hirata, *Tetrahedron Lett.*, 1986, **27**, 2113.
- 162 W. D. Schmitz, N. B. Messerschmidt and D. Romo, *J. Org. Chem.*, 1998, **63**, 2058.
- 163 L. Murray, H. Hamit, J. N. A. Hooper, L. Hobbs and R. J. Capon, *Aust. J. Chem.*, 1995, **48**, 1899.
- 164 L. Murray, A. T. R. Sim, J. A. P. Rostas and R. J. Capon, *Aust. J. Chem.*, 1993, **46**, 1291.
- 165 S. Urban and R. J. Capon, *Aust. J. Chem.*, 1992, **45**, 1255.
- 166 Y. Liu, B. H. Bae, N. Alam, J. Hong, C. J. Sim, C. Lee, K. S. Im and J. H. Jung, *J. Nat. Prod.*, 2001, **64**, 1301.
- 167 K. Choi, J. Hong, C.-O. Lee, D. Kim, C. J. Sim, K. S. Im and J. H. Jung, *J. Nat. Prod.*, 2004, **67**, 1186.
- 168 G. Cimino, S. de Stefano, L. Minale and E. Fattorusso, *Tetrahedron*, 1972, **28**, 333.
- 169 D. J. Faulkner, *Tetrahedron Lett.*, 1973, **14**, 3821.
- 170 I. Rothberg and P. Shubiak, *Tetrahedron Lett.*, 1975, **16**, 769.
- 171 F. Cafieri, E. Fattorusso, C. Santacroce and L. Minale, *Tetrahedron*, 1972, **28**, 1579.
- 172 C. J. Barrow, J. W. Blunt, M. H. G. Munro and N. B. Perry, *J. Nat. Prod.*, 1988, **51**, 275.
- 173 M. Kernan, R. C. Cambie and P. R. Bergquist, *J. Nat. Prod.*, 1991, **54**, 265.
- 174 C. J. Barrow, J. W. Blunt, M. H. G. Munro and N. B. Perry, *J. Nat. Prod.*, 1988, **51**, 1294.
- 175 C. J. Barrow, J. W. Blunt and M. H. G. Munro, *J. Nat. Prod.*, 1989, **52**, 346.
- 176 M. R. Kernan and D. J. Faulkner, *J. Org. Chem.*, 1988, **53**, 2773.
- 177 M. Ishibashi, M. Kurosaki, Y. Mikami and J. Kobayashi, *Nat. Prod. Lett.*, 1993, **3**, 189.
- 178 R. J. Capon, T. R. Dargaville and R. Davis, *Nat. Prod. Lett.*, 1994, **4**, 51.
- 179 A. Martinez, C. Duque, N. Sato, R. Tanaka and Y. Fujimoto, *Nat. Prod. Lett.*, 1995, **6**, 1.
- 180 R. Davis and R. J. Capon, *Aust. J. Chem.*, 1994, **47**, 933.
- 181 A. Martinez, C. Duque, N. Sato and Y. Fujimoto, *Chem. Pharm. Bull.*, 1997, **45**, 181.
- 182 U. Höller, G. M. König and A. D. Wright, *J. Nat. Prod.*, 1997, **60**, 832.
- 183 K. Takabe, H. Hashimoto, H. Sugimoto, M. Nomoto and H. Yoda, *Tetrahedron: Asymmetry*, 2004, **15**, 909.
- 184 Y. H. Jiang, S.-H. Ryu, E.-Y. Ahn, S. You, B.-J. Lee, J. H. Jung and D.-K. Kim, *Nat. Prod. Sci.*, 2004, **10**, 272.
- 185 K. McPhail, M. T. Davies-Coleman and P. Coetzee, *J. Nat. Prod.*, 1998, **61**, 961.
- 186 R. P. Walker, J. E. Thompson and D. J. Faulkner, *J. Org. Chem.*, 1980, **45**, 4976.
- 187 Y. Noda, H. Hashimoto and T. Norizuki, *Heterocycles*, 2001, **55**, 1839.
- 188 E. Fattorusso, V. Lanzotti, S. Magno, L. Mayol, M. DiRosa and A. Ialenti, *Bioorg. Med. Chem. Lett.*, 1991, **1**, 639.
- 189 M.-L. Bourguet-Kondracki, C. Debitus and M. Guyot, *J. Chem. Res. (S)*, 1996, 192.
- 190 S. De Rosa, A. De Giulio, A. Crispino, C. Iodice and G. Tommonaro, *Nat. Prod. Lett.*, 1997, **10**, 267.
- 191 G. Cimino, S. De Stefano and L. Minale, *Tetrahedron*, 1972, **28**, 1315.
- 192 M. Tsoukatou, H. Siapi, C. Vagias and V. Roussis, *J. Nat. Prod.*, 2003, **66**, 444.
- 193 M. S. Butler and R. J. Capon, *Aust. J. Chem.*, 1992, **45**, 1705.
- 194 S. De Rosa, A. Milone, A. De Giulio, A. Crispino and C. Iodice, *Nat. Prod. Lett.*, 1996, **8**, 245.
- 195 M. R. Kernan and D. J. Faulkner, *J. Org. Chem.*, 1988, **53**, 4574.
- 196 S. De Rosa, A. De Giulio, A. Crispino, C. Iodice and G. Tommonaro, *Nat. Prod. Lett.*, 1997, **10**, 7.
- 197 A. Martinez, C. Dunque, N. Hara and Y. Fujimoto, *Nat. Prod. Lett.*, 1995, **6**, 281.
- 198 A. Martinez, C. Duque and Y. Fujimoto, *Lipids*, 1997, **32**, 565.
- 199 M. B. Yunker and P. J. Scheuer, *J. Am. Chem. Soc.*, 1978, **100**, 307.
- 200 E. D. de Silva and P. J. Scheuer, *Tetrahedron Lett.*, 1980, **21**, 1611.
- 201 F. W. Sum and L. Weiler, *J. Am. Chem. Soc.*, 1979, **101**, 4401.
- 202 H. Bousserouel, M. Litaudon, B. Morleo, M.-T. Martin, O. Thoison, O. Nosjean, J. A. Boutin, P. Renard and T. Sevenet, *Tetrahedron*, 2005, **61**, 845.
- 203 G. Bifulco, I. Bruno, L. Minale, R. Riccio, C. Debitus, G. Bourdy, A. Vassas and J. Lavayre, *J. Nat. Prod.*, 1995, **58**, 1444.
- 204 S. Kato, K. Shindo, H. Kawai, A. Odagawa, M. Matsuoka and J. Mochizuki, *J. Antibiot.*, 1993, **46**, 892.
- 205 V. R. Macheria, J. Liu, C. Bellows, S. Teisan, B. Nicholson, K. S. Lam and B. C. M. Potts, *J. Nat. Prod.*, 2005, **68**, 780.
- 206 K. A. E. Sayed, A. M. S. Mayer, M. Kelly and M. T. Hamann, *J. Org. Chem.*, 1999, **64**, 9258.